CHI Learning & Development System (CHILD)

**Project Title** 

Time is Brain

**Project Lead and Members** 

• Dr Deborah Khoo

• Dr Prakash Rameshchandra Paliwal

• Dr Anil Gopinathan

**Organisation(s) Involved** 

National University Health System

**Project Category** 

**Clinical Improvement** 

**Keywords** 

Clinical Improvement, Process Redesign, Quality Improvement, Workflow

Improvement, Acute Stroke, Efficient Care, Timely Intervention, Early Management,

Endovascular Treatment, Timely Anesthesia Delivery, Anaesthesia, Facilitated

Conscious Sedation, Cost Saving, Neurology, Neuro-Stroke Department, Quality

Improvement Methodology, National University Hospital, Fish Bone Diagram, Pareto

Chart

Name and Email of Project Contact Person(s)

Name: Dr Deborah Khoo, Anaesthesia, National University Hospital

Email: deborah\_khoo@nuhs.edu.sg



# Time is Brain

Dr. Deborah Khoo

Dr. Prakash Rameshchandra Paliwal

Dr. Anil Gopinathan

National University Hospital



# INTRODUCTION

### When a Stroke happens, the clock starts ticking...

**Pre-hospital** Variable length of time from Paramedics activate stroke onset

Difficult to modify, not within our control. May be hours

### **Arrival in ED**

standby for stroke. Immediate CT brain, diagnosis and stroke team activation

Modifiable – has been optimized in a previous CPIP project. Average <10 min

#### **Treatment**

1)tPA 2) **EVT** 

Modifiable – time to EVT can be optimized Average 144 min to door to needle (2015 data)

### TIME IS BRAIN!

"Reperfusion should be achieved as early as possible, and when treatment is initiated beyond 6 hours from symptom onset, the effectiveness of endovascular therapy is uncertain." - 2015 AHA/ASA Focused Update of the 2013 Guidelines for the Early Management of Patients With Acute Ischemic Stroke Regarding Endovascular Treatment. Stroke. 2015, June 29, 2015

## Define the problem:


What can anaesthesia do to make a difference to timing?


- Anaesthetist needed to administer facilitated conscious sedation, enable faster conversion to GA if required.
- Time is brain being ready to deliver the anaesthesia rapidly when the patient arrives can help reduce overall time.
- Hurdles: Need time to get manpower and move equipment to angiosuite, screen patient, set lines etc. - can take up to 30 minutes. Could be minimized if everything is ready when the patient arrives.
- 30 minutes is clinically significant! Translates to improvements in patient's neurological recovery and healthcare burden/costs.

# METHODOLOGY

### 1. Stakeholder team identifies problems.


#### 2. Pareto Charting







#### 3. Implementation of solutions

| PROBLEM  | INTERVENTION   | DATE OF IMPLEMENTATION |  |
|--|--|------------------------|--|
| No unified guidelines/pathway                                  | <ul> <li>Problem identified</li> <li>Stakeholders gathered</li> <li>Formulated, reviewed and agreed on workflow.</li> <li>Workflow presented to relevant departments prior to implementation.</li> </ul> | 9 May 2016             |  |
| Neurology not familiar with anaesthesia workflow and processes |  |                        |  |
| Not clear who from anaesthesia to call                         |  |                        |  |
| Neurostroke activation pathway not understood by anaesthetist  |  |                        |  |
| Anaesthesia equipment not in DDI                               | DDI agreeable for<br>Anaesthesia machine placed<br>into DDI room   | 8 Nov 2016             |  |



# RESULTS





#### **Cost Savings – Implementation of** Workflow

|  | Pre-intervention      | Post-intervention      |
|--|-----------------------|------------------------|
| Mean Length of Acute Inpatient Stay  | 18 days               | 14 days                |
| Modified Rankin Scale for Neurologic Disability 0 = no symptoms at all 1 = no significant disability 2 = slight disability 3 = moderate disability 4 = moderately severe disability 5 = severe disability 6 = dead | 3                     | 3                      |
| Note:  • Door to anaesthesia time is only  | one of many contribut | ing factors. This is a |

continuous team effort, led by the Neuro-stroke department, and greatly helped by the expertise of the Interventional Radiologist. Length of stay and patient outcomes are impacted by many other factors, e.g. pre-existing comorbidities, age of patient.

"This project put in place a system that allowed everyone to be ready within minutes of activation. When I witnessed for myself how a patient improved tremendously after a clot retrieval, I realised how big an impact a few crucial minutes and our teamwork could make on the patient's outcome." -Ms. Mary Rose Gomez Calderon, Assistant Nurse Clinician, Anaesthesia Unit Operating Theatre.

# CONCLUSION

- EVT caseload is increasing year on year, with growing international interest and data.
- This project helps us be future ready and better able to cope with caseload.
- Constant reassessment, mutual avenues for feedback and suggestions for improvement when needed.
- Celebrate improvements and good patient outcomes!
- Positive feedback and encouragement to individuals.

**ACKNOWLEDGEMENTS:** Our grateful thanks to Dr Wong WH, Dr Teoh HL, A/Prof R Seet and our CPIP supervisors for their guidance and support and for believing in us.